

Assessing Invertebrate epifaunal habitat preference in a shallow coastal bay

Lowell Andrew Iporac*, Manny Vera, Ligia Collado-vides

Department of Biological Sciences, Florida International University



Background Information

- Macroalgae are known to contribute to habitat complexity in shallow coastal systems; habitat enhancement is dependent on the specific macroalgal species (Norkko et al. 2000) (Fig. 1).
- Biscayne Bay, localized close to a metropolitan area in South Florida, undergoes heavy management and restoration activities impacting adjacent coastal communities (Morrison, 2015). Previous studies show that freshwater pulses in the area cause fluctuations of salinity and temperature, leading to changes in macroalgal species composition and invertebrate epifaunal distributions (Alleman et al. 2013, Brooks 1982, Charkhian 2014, Collado-Vides et al. 2011)
- Previous studies on other Floridian coastal areas, such as the Indian River Lagoon and Tampa Bay, showed that epifaunal species abundance varied between the drift algae and seagrass beds, yet have similar species composition between habitat types. However, the dominance of specific epifaunal species between habitats differ between coastal areas (Knowles and Bell, 1994, Virnstein and Howard 1982).
- Here we test if invertebrate epifauna display habitat preference within Deering Estate at Biscayne Bay. Understanding epifaunal habitat preference between macrophyte habitats provides insight in how coastal communities function, as well as provide a baseline to evaluate environmental disturbances such as potential consequences of water management strategies.
- We hypothesize that differences in epifaunal communities exist due to structural differences between macrophyte habitats.



Figure 1: A red macroalgal mat embedded within a seagrass bed

Objectives

- Characterize the red macroalgal mat (RAM) and benthic seagrass (BSG) habitats based on macrophyte species composition.
- Determine if there is habitat preference among invertebrate epifaunal groups between the two habitat types.
- Determine if epifaunal community distributions is based more on habitat selection or environmental factors.

Methods

- Sampling was conducted at four different sites in Deering Estate, Biscayne Bay once every two to three months (October 2017, December 2017, and March 2018) (Fig 2)



Figure 2: Study location (Deering Estate) and sites, relative to geographical location

- Five samples of BSG and RAM habitats were collected per site by encircling a plastic bag over the macrophyte habitat to ensure capture of invertebrate epifauna.
- Once samples were returned to the lab, all macrophyte species per sample are separated based on species or genus (table 1), and are weighed for wet and dry biomass.
- Invertebrate epifauna within that same sample were sorted into broad taxonomic levels (table 2) and were counted for abundance.
- Historical salinity data from previous monitoring efforts (2015-2017) were collected using a YSI and current salinity data (2018 onward) were collected using a refractometer.
- Wilcoxon Rank Sum tests were used to compare epifaunal species richness and total macrophyte biomass between habitats. Student's t-test were used to compare log-transformed total epifaunal abundances between habitats
- Multivariate analyses (PERMANOVA) were used to compare epifaunal and macrophyte species composition between three factors (habitat, sampling date, and site).

Table 1: Species list of macrophytes between habitats

Species	Abbr.	Phylum	Habitat			
			RAM		BSG	
			Present?	Avg. Wet Biomass	Present?	Avg. Wet Biomass
<i>Thalassia testudinum</i>	Thal	Tracheophyta	X	5.139	X	19.836
<i>Halodule wrightii</i>	Halo	Tracheophyta	X	1.268	X	1.759
Diatoms	Diat	Bacillariophyta	X	0.005	X	0.878
<i>Penicillus capitatus</i>	Penc	Chlorophyta	X	0.026	X	0.032
<i>Batophora occidentalis</i>	Bato	Chlorophyta	X	0.529		
<i>Anadyomene stellata</i>	Anad	Chlorophyta	X	0.085	X	0.000056
<i>Digenea simplex</i>	Dige	Rhodophyta	X	6.866	X	0.00349
<i>Chondria sp.</i>	Chond	Rhodophyta	X	4.941	X	0.201
<i>Laurencia sp.</i>	Laur	Rhodophyta	X	15.739	X	0.283
<i>Spyridia filamentosa</i>	Spyr	Rhodophyta	X	5.657	X	0.00356
<i>Acanthophora spicifera</i>	Acan	Rhodophyta	X	1.504		
<i>Polysiphonia sp.</i>	Plys	Rhodophyta	X	0.147	X	0.04
<i>Ceramium sp.</i>	Cera	Rhodophyta	X	0.006	X	0.000056
<i>Jania sp.</i>	Jani	Rhodophyta	X	0.11	X	0.00005
<i>Centroceras sp.</i>	Cntr	Rhodophyta	X	0.021		
<i>Sargassum sp.</i>	Srgs	Ochrophyta	X	0.004		
Misc. leaves and Bark	LvsBk	Tracheophyta	X	0.272		

Total macrophyte biomass by habitat




Figure 3: Comparison of total macrophyte wet biomass between habitat

Figure 4: Comparison of Historical Salinity data between sites (left) and months (right) - (compiled from sampling 2015-2018)




Table 2: List of invertebrate epifauna between habitats

Group	Taxon	Abbr.	Habitat			
			RAM		BSG	
			Rel. Freq.	Total Abundance	Rel. Freq.	Total Abundance
Amphipod	Amphipoda	Amph	0.97	2670	0.89	239
Isopod	Isopods	Isop	0.74	378	0.22	6
Tanaid	Tanaidacea	Tana	0.18	34	0.28	77
Bivalve	Bivalva	Biva	0.97	5162	0.72	86
Gastropod	Gastropoda	Gast	1.00	3024	0.94	157
Chiton	Polyplacophora	Chit	0.11	4	0.22	10
Caridean Shrimp	Caridea	Shri	0.29	17	0.17	3
Hermit Crab	Paguroidea	Herm	0.42	98	0.17	7
Crabs	Brachyura	Crab	0.32	25	0.00	0
Starfish	Asteroidea	Aste	0.08	4	0.00	0
Polychaete Worms	Polychaeta	Polyc	0.79	179	0.78	246
Insect	Insecta	Inse	0.08	14	0.00	0
Ostracod	Ostracoda	Ost	0.05	6	0.06	8
Sipunculid Worms	Sipuncula	Sipu	0.05	5	0.11	2
Total				11620		841

Figure 5: Relative frequencies of epifaunal groups between habitats




Figure 6: Comparison of total invertebrate abundance (left) and species richness (right) between habitat types




Figure 7: Macrophyte wet biomass nMDS Multivariate analyses





Figure 8: Epifaunal abundance nMDS Multivariate analyses



Conclusion

- While many macrophyte species exist in both habitats, there are differences in the dominance of specific macrophyte species between habitats.
- BSG habitats are dominated by *Thalassia testudinum* and *Halodule wrightii*, while RAM habitats are dominated by rhodophytes such as *Laurencia sp.* and *Digenea simplex*
- RAM Habitats show higher species richness, relative frequency, and abundance of epifauna compared to BSG habitats.
- Epifaunal species composition seem to not only be determined by habitat choice, but also by other environmental factors that could arise from seasonality (date) and site-specific conditions.
- Differences in salinity exist between months, but not between sites, suggesting salinity may not be one factor contributing to invertebrate habitat preference,

Acknowledgements

This research was supported by the Broward Shell Club of West Palm Beach FL, and the FIU Center of Research and Excellence in Science and Technology - Center of Aquatic Chemistry and Environment (FIU CREST-CACHE).

References

- Alleman, R., Stabenau, E., Charkhian, B. & Brown, R. 2013. Pilot project tests for supplemental water deliveries to Biscayne Bay: After action assessment.
- Brook, I.M. 1982. The effect of freshwater canal discharge on the stability of two seagrass benthic communities in Biscayne National Park, Florida. *Oceanologica Acta, Special Issue*.
- Charkhian, B. 2014. Biscayne Bay coastal wetlands restoration benefit.
- Collado-Vides, L., Mazzei, V., Thyberg, T. & Lirman, D. 2011. Spatio-temporal patterns and nutrient status of macroalgae in a heavily managed region of Biscayne Bay, Florida, USA. *Botanica Marina*. 54:377-90.
- Knowles, L.L. & Bell, S.S. 1998. The Influence of Habitat Structure in Faunal-Habitat Associations in a Tampa Bay Seagrass System, Florida. *Bulletin of Marine Science*. 62:781-94.
- Morrison, M. 2015. South Florida water management district - Biscayne Bay coastal wetlands project. US Department of the Interior and US Army Corps of Engineers.
- Norkko, J., Bonsdorff, E. & Norkko, A. 2000. Drifting algal mats as an alternative habitat for benthic invertebrates: Species specific responses to a transient resource. *Journal of Experimental Marine Biology and Ecology*. 248:79-104.